



Systems biology

Breeze: an integrated quality control and data analysis application for high-throughput drug screening

Swapnil Potdar ^{1,*}, Aleksandr Ianevski^{1,2}, John-Patrick Mpindi¹, Dmitrii Bychkov¹, Clément Fiere¹, Philipp Ianevski¹, Bhagwan Yadav¹, Krister Wennerberg^{1,3}, Tero Aittokallio ^{1,2,4}, Olli Kallioniemi^{1,5}, Jani Saarela^{1,†} and Päivi Östling^{1,5,†}

¹Institute for Molecular Medicine Finland (FIMM), University of Helsinki, FI-00290 Helsinki, Finland, ²Department of Computer Science, Helsinki Institute for Information Technology (HIIT), Aalto University, FI-02150 Espoo, Finland, ³Biotech Research & Innovation Centre, University of Copenhagen, 2200 Copenhagen N, Denmark, ⁴Department of Mathematics and Statistics, University of Turku, FI-20014 Turku, Finland and ⁵Science for Life Laboratory (SciLifeLab), Department of Oncology and Pathology, Karolinska Institutet, 171 65 Solna, Stockholm, Sweden

*To whom correspondence should be addressed.

†The authors wish it to be known that, in their opinion, the last two authors should be regarded as Joint Last Authors.

Associate Editor: Alfonso Valencia

Received on October 11, 2019; revised on January 27, 2020; editorial decision on February 20, 2020; accepted on February 26, 2020

Abstract

Summary: High-throughput screening (HTS) enables systematic testing of thousands of chemical compounds for potential use as investigational and therapeutic agents. HTS experiments are often conducted in multi-well plates that inherently bear technical and experimental sources of error. Thus, HTS data processing requires the use of robust quality control procedures before analysis and interpretation. Here, we have implemented an open-source analysis application, Breeze, an integrated quality control and data analysis application for HTS data. Furthermore, Breeze enables a reliable way to identify individual drug sensitivity and resistance patterns in cell lines or patient-derived samples for functional precision medicine applications. The Breeze application provides a complete solution for data quality assessment, dose–response curve fitting and quantification of the drug responses along with interactive visualization of the results.

Availability and implementation: The Breeze application with video tutorial and technical documentation is accessible at <https://breeze.fimm.fi>; the R source code is publicly available at <https://github.com/potdarswapnil/Breeze> under GNU General Public License v3.0.


Contact: swapnil.potdar@helsinki.fi

Supplementary information: [Supplementary data](#) are available at *Bioinformatics* online.

1 Introduction

Advances in automated liquid dispensing, assay miniaturization, systems biology as well as *ex vivo* disease models have boosted the development of high-throughput cell-based functional testing platforms. Thousands of conventional cell lines have been screened with hundreds of compounds in several large-scale projects, such as the Cancer Cell Line Encyclopedia (Barretina *et al.*, 2012), Genomics of Drug Sensitivity in Cancer (Yang *et al.*, 2013), Cancer Therapeutics Response Portal (Seashore-Ludlow *et al.*, 2015) and Genentech Cell Line Screening Initiative (Haverty *et al.*, 2016). Similar drug testing efforts have been applied on primary cell models to generate individualized drug profiles for drug repurposing, patient stratification and for the identification of potential drug combinations (Kodack *et al.*, 2017; Lee *et al.*, 2018; Pemovska *et al.*,

2013, 2015; Saeed *et al.*, 2017). One common end-point in cell-based drug testing is cell viability and/or toxicity readouts generated over multiple concentrations in microwell plates (96-, 384- and 1536-well formats), where plate layout and placement of controls play an important role to minimize the risk of experimental errors influencing data quality (Mpindi *et al.*, 2015). Therefore, quality control (QC) process is required to ensure compliant and reproducible drug testing readouts from the assay. The dose–response curve-fitting process enables direct translation of the raw cell viability measurements based on several intensity scoring technologies to clinically interpretable dose values. The fitted dose–response curve is then used to summarize and quantify the observed response into a single metric, such as IC₅₀ and EC₅₀ dose or as an absolute area under the curve (AUC) or drug sensitivity score (DSS; Yadav *et al.*, 2015). To date, there are several, freely available analysis tools



functionality in terms of QC, summary metrics and visualization of different aspects of HTS experiments.

Acknowledgements

We would like to thank all FIMM Breeze application users for valuable input and beta testing.

Funding

The FIMM High-Throughput Biomedicine Unit is financially supported by the University of Helsinki and Biocenter Finland. The work has received funding from the Sigrid Juselius Foundation, Cancer Society of Finland, Academy of Finland [292611, 326238 and 310507 to T.A.], the Magnus Ehrnrooth Foundation, Biocentrum Helsinki and St. Baldrick's Foundation.

Conflict of Interest: none declared.

References

- Barretina, J. *et al.* (2012) The Cancer Cell Line Encyclopedia enables predictive modelling of anticancer drug sensitivity. *Nature*, **483**, 603–607.
- Boutros, M. *et al.* (2006) Analysis of cell-based RNAi screens. *Genome Biol.*, **7**, R66.
- Findlay, J.W.A. and Dillard, R.F. (2007) Appropriate calibration curve fitting in ligand binding assays. *AAPS J.*, **9**, E260–E267.
- Chen, L. *et al.* (2016) mQC: a heuristic quality-control metric for high-throughput drug combination screening. *Sci. Rep.*, **6**, 37741.
- Fourches, D. *et al.* (2014) HTS navigator: freely accessible cheminformatics software for analyzing high-throughput screening data. *Bioinformatics*, **30**, 588–589.
- Haverty, P.M. *et al.* (2016) Reproducible pharmacogenomic profiling of cancer cell line panels. *Nature*, **533**, 333–337.
- Kodack, D.P. *et al.* (2017) Primary patient-derived cancer cells and their potential for personalized cancer patient care. *Cell Rep.*, **21**, 3298–3309.
- Lee, S.H. *et al.* (2018) Tumor evolution and drug response in patient-derived organoid models of bladder cancer. *Cell*, **173**, 515–528.e517.
- Mpindi, J.P. *et al.* (2015) Impact of normalization methods on high-throughput screening data with high hit rates and drug testing with dose-response data. *Bioinformatics*, **31**, 3815–3821.
- Mpindi, J.P. *et al.* (2016) Consistency in drug response profiling. *Nature*, **540**, E5–E6.
- Pemovska, T. *et al.* (2015) Axitinib effectively inhibits BCR-ABL1(T315I) with a distinct binding conformation. *Nature*, **519**, 102–105.
- Pemovska, T. *et al.* (2013) Individualized systems medicine strategy to tailor treatments for patients with chemorefractory acute myeloid leukemia. *Cancer Discov.*, **3**, 1416–1429.
- Saeed, K. *et al.* (2017) Comprehensive drug testing of patient-derived conditionally reprogrammed cells from castration-resistant prostate cancer. *Eur. Urol.*, **71**, 319–327.
- Seashore-Ludlow, B. *et al.* (2015) Harnessing connectivity in a large-scale small-molecule sensitivity dataset. *Cancer Discov.*, **5**, 1210–1223.
- Smirnov, P. *et al.* (2016) PharmacGx: an R package for analysis of large pharmacogenomic datasets. *Bioinformatics*, **32**, 1244–1246.
- Vølund, A. (1978) Application of the four-parameter logistic model to bioassay: comparison with slope ratio and parallel line models. *Biometrics*, **34**, 357–365.
- Yadav, B. *et al.* (2015) Quantitative scoring of differential drug sensitivity for individually optimized anticancer therapies. *Sci. Rep.*, **4**, 5193.
- Yang, W. *et al.* (2013) Genomics of Drug Sensitivity in Cancer (GDSC): a resource for therapeutic biomarker discovery in cancer cells. *Nucleic Acids Res.*, **41** (Database issue), D955–D961.
- Zhang, J.H. (1999) A simple statistical parameter for use in evaluation and validation of high throughput screening assays. *J. Biomol. Screen.*, **4**, 67–73.
- Zhang, X.D. (2007) A pair of new statistical parameters for quality control in RNA interference high-throughput screening assays. *Genomics*, **89**, 552–561.